



**NADORCOTT MANAGED WITH OUR  
BIOTECHNOLOGY 09/01/2020**

## BIOTECHNOLOGICAL MEASURES TO OBTAIN A PREMIUM PRODUCTION:

- Achieve a **MIXED SPROUT** with the ideal number of flowers and control fungal diseases.
- Maintain an **OPTIMAL ROOT SYSTEM** with an excellent assimilation of nutrients and control the electrical conductivity without exceeding the levels of 2.2 in the bulb.
- Use the **HORMONE-FREE FRUIT THINNING AND LIGHTENING** treatment at the beginning of the physiological fall when fruits fall with calyx and make 2 to 3 applications 8 days apart.



## TO CONTROL FUNGAL DISEASES AND ACHIEVE A BALANCE IN THE CROP

### STIMULATION TREATMENT

Conduction: from February 15th to March 10th

*Although the fruit is hanging, the treatment is carried out, it is Zero Residue and without side effects for the fruit*



**CUPRO-PLUS MAX** \_\_\_\_\_ 3 L. x 1000 L.



**ALGATRON** \_\_\_\_\_ 1,25 L. x 1000 L.



**MELASTOP** \_\_\_\_\_ 1,5 L. x 1000 L.





Minimum volume 4.000 L./ha.

## **TO ACHIEVE A MIXED SPROUT WITH FEW FLOWERS AND CONTROL FUNGAL DISEASES**

### **STIMULATION TREATMENT**

Conduction: mid or second half of March

*Although the fruit is hanging, the treatment is carried out, it is Zero Residue and without side effects for the fruit*

	<b>ENDOFOSFITO</b> _____ 5	L. x 1000 L.
	<b>AMINOPLUS</b> _____ 2,5	L. x 1000 L.
	<b>EDYMAG</b> _____ 2,5	L. x 1000 L.
	<b>MELASTOP</b> _____ 1,5	L. x 1000 L.

Minimum volume 2.500 L./ha.

## TO ACHIEVE A MIXED SPROUT WITH FEW FLOWERS AND WELL CONFIGURED AND BALANCED OVARIES

### STIMULATION TREATMENT

Conduction: early April

*Although the fruit is hanging, the treatment is carried out, it is Zero Residue and without side effects for the fruit*



**AMINOPLUS**\_\_\_\_\_2,5 L. x 1000 L.



**HYDROCAL**\_\_\_\_\_2,5 L. x 1000 L.



**ZIMA-L MAX**\_\_\_\_\_2,5 L. x 1000 L.



**MELASTOP**\_\_\_\_\_1,5 L. x 1000 L.

Minimum volume 2.500 L./ha.

**TO MAINTAIN AN OPTIMAL ROOT SYSTEM WITH EXCELLENT ASSIMILATION OF  
NUTRIENTS AND CONTROL E.C. WITHOUT EXCEEDING THE LEVELS OF 2.2 IN  
THE BULB**

**ROOT TREATMENT**

**1ª Application MARCH:**



**COUPÉ REGENERACIÓN PLUS** \_\_\_\_\_ 80 L./ha.

**Once the physiological fall is over, at the start of fattening:**



**COUPÉ REGENERACIÓN PLUS** \_\_\_\_\_ 80 L./ha.

**In times of overexertion during the summer:** additional applications of 60 to 80 L./ha.,  
Applied twice 12-14 days apart improve fattening significantly.

**EDYFOS** is a **powerful source of assimilable phosphorus** (45% w/w) mainly intended to facilitate the assimilation of this element in adverse circumstances. Phosphorus is an element, par excellence, of quality. **Molybdenum** (0.5% w/w) provided by **EDYFOS** increases the level of pollen fertility.

### ROOT TREATMENT

### DURING MARCH AND APRIL



**EDYFOS** \_\_\_\_\_ 10 L./ha. and application.

Make between 3 and 4 applications.







**HIGH PRODUCTION VARIETIES  
(NADORCOTT, TANGO, ESBAL,  
SANGUINELLI)**

**USE THE FRUIT LIGHTENING TREATMENT WITHOUT HORMONES AT THE ONSET OF  
THE PHYSIOLOGICAL FALL WHEN FRUIT FALLS WITH CHALICE**

**FREE HORMONE FRUIT LIGHTENING TREATMENT**

Conduction: from 2 to 3 application 8 days apart

	<b>ENDOFOSTEC</b> _____	4	L. x 1000 L.
	<b>NITROPLUS</b> _____	4	L. x 1000 L.
	<b>MAX QUALITY</b> _____	3	L. x 1000 L.
	<b>HYDROCAL</b> _____	3	L. x 1000 L.
	<b>ZIMA-L MAX</b> _____	2,5	L. x 1000 L.
	<b>MELASTOP</b> _____	1,5	L. x 1000 L.



**SUPPORT THIS TREATMENT WITH ORGANOVITAL PLUS APPLICATIONS WHEN THE  
1ST LIGHTENING TREATMENT IS DONE**

**ROOT TREATMENT**



**ORGANOVITAL PLUS** \_\_\_\_\_ 100 L. x ha.



**CUPRO-PLUS MAX** is a new and powerful formulation of **organic copper** with high assimilation and **easy mobility in the vegetative mass** designed to recover the copper levels in vegetables, keeping the vital and nutritional functions of the crop in balance.

**CONTROL OF CRYPTOGAMIC DISEASES TREATMENT**

Carry out 3 annual applications:

1st application: from February 15th to March 31st

2nd application: beginning of summer

3rd application: end of summer



**CUPRO-PLUS MAX** \_\_\_\_\_ 2,5      L. x 1000 L.



**MELASTOP** \_\_\_\_\_ 1,5      L. x 1000 L.

Minimum volume 3.500 L./ha.

**ENDOFOSTEC** has a **vasodilator/stimulating effect** and it is recommended to apply when it is necessary to recover the vegetative rhythm of the crop that has fallen due to root asphyxia or overexertion. It is an excellent tool to increase fattening actions.

**ENDOFOSTEC** is a great tool to avoid the loss of fruits caused by rain.

### **ENDOPARASITE DISEASES CONTROL TREATMENT**

Carry out 3 annual applications:

1st application: during the spring

2nd application: end of summer

3rd application: beginning of autumn



**ENDOFOSTEC** \_\_\_\_\_ 4 L. x 1000 L.



**MELASTOP** \_\_\_\_\_ 1,5 L. x 1000 L.

Minimum volume 3.000 L./ha.