



EDYPRO
biotecnologia



**Increase yield and improve the
quality of POME FRUITS with the use
of BIOTECHNOLOGY**



Treatment for soil preparation Assimilation of nutrients from the soil

Radicular Application

COUPÉ REGENERACIÓN Plus

1st application: from 50 to 80 L/ha., in the early spring.

At the beginning of fruit set, maturation and moments of overexertion: applications of 60 to 80 L./ha, improve maturation significantly.



Treatment for unbalanced bloom and excess production

Foliar Application

We recommend 2 applications in the spring of **ALGATRON** associated with **MELASTOP** with 50% of open flower and repeat to 20 days:

ALGATRON

4 L. x 1000 L.

MELASTOP

1,5 L. x 1000 L.

With minimum consumption of 500 to 600 L. / ha of broth.

Note: This combination is compatible with the usual plant protection treatments.



Treatment for Maturation

Foliar Application

MAX QUALITY

4 L. x 1000 L.

CA-FRUIT

4 L. x 1000 L.

ZIMA-L MAX

2,5 L. x 1000 L.

MELASTOP

1,5 L. x 1000 L.

1st application: before the seed's hardening.

2nd application: After 15 days of the 1st application.

NOTE. With minimum consumption of 500 L / ha. of broth.



Treatment for Rippening

Foliar Application

1st application make 25 days of the desired collection date:

MADUR FRUIT FOLIAR

5 L. x 1000 L.

MELASTOP

1,5 L. x 1000 L.

2nd application 10 days after the 1st treatment:

MADUR FRUIT FOLIAR

5 L. x 1000 L.

MELASTOP

1,5 L. x 1000 L.

NOTE. With minimum consumption of 600 L. / ha. of broth.

