



**EDYPRO**  
biotecnologia



**Increase the yield and improve the  
quality of Persimon with the use of  
BIOTECHNOLOGY**



# STIMULATION OF SPROUTING AND VEGETATIVE DEVELOPMENT

## Radicular Application

**COUPÉ REGENERACIÓN Plus** 40 L. / ha.

**FERROMAN 5%** 10 kg. / ha.

**FER MAGNESIUM** 28-42 kg./ ha.

**Moment of the application:** April - May

Apply each product separately and distance 10-15 days.

**NOTE:** Discontinue fertilizer when applying  
**COUPÉ REGENERACIÓN Plus** and **Ferroman 5%**.



# **TREATMENTS FOR OPTIMUM VEGETATIVE DEVELOPMENT AND CONTROL OF MYCOSPHAERELLA**

**Moment of the applications: starting in April.**

**THREE CONSECUTIVE TREATMENTS  
EVERY 15-20 DAYS**



# **1st TREATMENT**

## **Foliar Application**

**ULTRAMAN**

**2,5 L. x 1000 L.**

**EDYMAG**

**2,5 L. x 1000 L.**

**MELASTOP**

**1,5 L. x 1000 L.**

**Fungicide For Mycosphaerella**

**NOTE.** For weakened plantations, it is recommended to add to this treatment:

**ALGATRON**

**2,5 L. x 1000 L.**



**EDYPRO**  
biotecnologia



## **2nd TREATMENT**

### **Foliar Application**

**ULTRAMAN**

**2,5 L. x 1000 L.**

**HYDROCAL**

**2,5 L. x 1000 L.**

**MELASTOP**

**1,5 L. x 1000 L.**

**Fungicide For Mycosphaerella**

**Moment of the application: 15 - 20 days after the 1<sup>st</sup> treatment.**



**EDYPRO**  
biotecnologia



## **3rd TREATMENT**

### **Foliar Application**

<b>ULTRAMAN</b>	<b>1,5 L. x 1000 L.</b>
<b>HYDROCAL</b>	<b>2,5 L. x 1000 L.</b>
<b>EDYMAG</b>	<b>2 L. x 1000 L.</b>
<b>MELASTOP</b>	<b>1,5 L. x 1000 L.</b>
<b>Fungicide For Mycosphaerella</b>	

**Moment of the application: 15-20 days after the 2<sup>nd</sup> treatment.**



# **CORRECTING CALCIUM DEFICIENCIES**

## **Radicular Application**

### **FER CALCIUM**

**Drip irrigation: 20 to 35 L. / ha. per application**

**Surface irrigation: 40 to 60 L. / ha. per application**

**NOTE:** Make 2-4 applications distanced from each other about  
15 days.



# **CORRECTING MAGNESIUM DEFICIENCIES**

## **Foliar Application:**

**EDYMAG** de 200 a 300 c.c./Ha.

## **Radicular Application:**

### **FER MAGNESIUM**

**Drip irrigation: 15 to 20 L. / ha. per application.**

**Surface irrigation: 25 to 60 L. / ha. per application.**





**EDYPRO**  
biotecnologia



## **CORRECTING MANGANESE DEFICIENCIES**

### **Foliar Application:**

**ULTRAMAN 200 to 300 c.c./100 L.**

Make minimum 2 applications per month.

### **Radicular Application:**

**ULTRAMAN 25 to 50 L. / ha.**

Subdivided in 2 or 3 applications.



## **TREATMENTS FOR MATURATION**

**Sometimes the evolution of the maturation is less than desired and therefore, we find little commercial sizes.**

**EDYPRO has developed a new system for radicular applications that allows increasing sizes and significantly reducing the % of small fruits.**



# TREATMENT FOR MATURATION

## Radicular Application

**SUPER ENGORDE CAQUI**

**15-25 L./ Ha. per application**

**1st application:** in August, choosing the time of application according to the need of the crop.

**2nd application:** first half of September or 20 to 25 days after the first application.

**Note:** *in the late varieties, an application can be made in the first half of October.*



# **SKIN PROTECTION TREATMENT**

## **Foliar Application**

**CA-FRUIT**

**3,5 L. x 1000 L.**

**MELASTOP**

**1,5 L. x 1000 L.**

Repeat every 12-14 days.

**NOTE:** if it is intended to green, add Gibberellic Acid.



**TAKE ADVANTAGE AND SUCCEED WITH US**