



Increase the production and improve the quality of Onion through BIOTECHNOLOGY



IMPROVING THE ONION PRODUCTION IN AUTUMN / WINTER CICLE

1st FASE: Rooting and start of Vegetative Development



BULB PREPARATION Radicular Application

Recommendation:

To achieve a shallow horizontal bulb without waterlogging (this situation, if successful, causes root asphyxia and encourages fungal attack).

EDYPRO HUMECTANTE 20 L. / ha.

Applied in a single irrigation dissolved from the first to the last drop. The duration of this irrigation will never exceed 2 hours.





ROOTING AND FAST DEVELOPMENT OF THE ROOT SYSTEM

Radicular Application

COUPÉ REGENERACIÓN Plus60 L. / ha.ENRAIZANTE ESPECIAL10 L. / ha.

Repeat in 12 days:

COUPÉ REGENERACIÓN Plus80 L. / ha.ENRAIZANTE ESPECIAL5 L. / ha.





Thrips problems in onion cultivation

REVOLUTION is a new generation product whose action through the serum effect is capable, if continuous applications are made, of controlling the thrips population and allowing the cultivation to be successfully carried out.

Foliar Application

REVOLUTION	5 L. x 1000 L.
MELASTOP	2 L. x 1000 L.

Treat at sunset and wet deeply. As it is a Zero Residue product, it can be applied 3 times a week without negative effects on the crop. It can be applied alone or in combination with other treatments.

It is compatible with the auxiliary fauna used to control thrips such as the Orius.





ROOTING AND RAPID DEVELOPMENT OF THE FOLIAR SYSTEM **Foliar Application HYDROCAL** 3 L. x 1000 L. **EDYFOS** 0,75 L. x 1000 L. BOTCAR 3 L. x 1000 L. **ENDOFOSTEC** 5 L. x 1000 L. 2 L. x 1000 L. MELASTOP

NOTE: Carry out 3 passes 10 days apart and add the chemical fungicide and/or insecticide that is considered.



IMPROVING THE ONION PRODUCTION IN AUTUMN / WINTER CICLE

2nd FASE: Vegetative Development





STEM DEVELOPMENT AND
VEGETATIVE RHYTHM
Foliar ApplicationHYDROCAL3 L. x 1000 L.ENDOFOSTEC or5 L. x 1000 L.

Associated with conventional fungicide treatments

NOTE. It is very important to look for the anti-stress effect caused by fungicides. For this we recommend accompanying the treatments with:

ALGATRON

MELASTOP

2 L. x 1000 L.

2 L. x 1000 L.



IMPROVING THE ONION PRODUCTION IN AUTUMN / WINTER CICLE

3rd FASE: Maturation





MATURATION PLAN AND RAPID BULB DEVELOPMENT

Radicular Application

ORGANOVITAL PLUS

60 L. / ha.

After 12 Days:

GROSKOLOR PLUS

80 L. / ha.





MATURATION PLAN AND RAPID BULB DEVELOPMENT

Foliar Application

EDYMAG	2,5 L. x 1000 L
MAX QUALITY	4 L. x 1000 L
CA-FRUIT	2,5 L. x 1000 L
ENDOFOSTEC	5 L. x 1000 L
MELASTOP	2 L. x 1000 L

NOTE: Add the fungicide and/or chemical insecticide considered.



Treated with MAX QUALITY